

**Ecology of migrant shorebirds in New Zealand, focussing
on Farewell Spit, North-West Nelson**

A thesis presented in partial fulfilment of the
requirements for the degree of Masterate in Ecology
at Massey University

Philip Frank Battley

1996

ABSTRACT

Migratory shorebirds are a dynamic component of New Zealand's coastal fauna, alternating between distant breeding and non-breeding grounds. The Red Knot *Calidris canutus*, Bar-tailed Godwit *Limosa lapponica* and Pied Oystercatcher *Haematopus ostralegus finschi* were studied on Farewell Spit, North-West Nelson. The first two species breed in the Arctic and migrate to New Zealand for the non-breeding season. Over the southern summer they experience low thermostatic costs and generally improving prey conditions over the summer. Oystercatchers are resident over autumn and winter, so experience rising costs and declining prey quality in some species. One bivalve species, *Macomona liliana*, shows seasonal depth changes in the sediment, and so is largely inaccessible even to a long-billed bird such as the oystercatcher.

Despite this, oystercatchers feed for less time than is available, and achieve intake rates sufficient to cover estimated needs. The energy needs of the Arctic waders rise as they prepare to migrate, and they achieve at least part of this by increasing the duration of feeding. Knots during spring tides in the premigratory period feed for the entire low-water period. Godwits are apparently less stressed, underutilising nocturnal feeding opportunities over summer. They are thought to increase feeding time by using this night-time feeding.

The high energy demands for migrating birds come from the need to deposit nutrients for migration, and knots around the Auckland region are estimated to increase in mass from 115 to 185 g prior to migration. Fat deposition is not the only physiological preparation, however, and a sample of knots shot from Northland (illegally, recovered by DoC) revealed complex interactions between organs. Large amounts of fat were deposited, mostly in a subcutaneous layer but also in the abdominal cavity. Muscle protein was also deposited in flight and heart muscles, presumably to prepare for the extreme effort involved in trans-oceanic flights. At the same time, digestive organs decreased in mass. This is interpreted as freeing up muscle protein which is then deposited in organs for use during flight.

Knots and godwits migrated from Farewell Spit in March. Most departures occurred in the evening and on rising tides. The former probably allows for the use of multiple navigational cues, while the latter may maximise feeding opportunities immediately before the flight. Most departures occurred after the passage of a low-pressure system or with the approach of a high-pressure system. This enabled favourable winds to be gained, so that the mean wind vector was a small tailwind. Thus, while departure directions were intermediate between the expected directions for flights to either Australia or northern New Zealand, it is probable they were able to fly across the

top of a high-pressure system and gain wind assistance for a direct flight to Australia. However, the variability in flight range estimates depending on assumptions of travel-speed and protein deposition makes predicting migration routes difficult.

Numbers of godwits have increased on Farewell Spit over the past decade, while oystercatchers have remained static. Knots have shown a slight decline. Knot numbers are independent of national census counts so are presumably determined largely by factors operating on Farewell Spit itself. A possible mechanism that could give to a slowly declining population could be if a certain sector of the population fails to deposit sufficient nutrients to successfully migrate and return. If site-fidelity is very high (as it generally is in waders) then a long-term decrease could ensue.

TABLE OF CONTENTS

	Title page	i
	Dedication	ii
	Abstract	iii
	Table of contents	v
	Acknowledgements	ix
	List of figures and tables	xii
Chapter 1	General introduction	1
Chapter 2	Study site: physical features, biological features and history	
	Geomorphology	8
	Fieldwork	9
	Climate	12
	Vegetation	12
	Fauna	13
	Status	14
	Scientific research	14
Chapter 3	Prey availability	
	Abstract	15
	Introduction	16
	Methods	18
	Results	
	Total biomass	20
	Size relationships in bivalves	25
	Sizes of shellfish	29
	Size-mass relationships	31
	Worms	40
	Discussion	42
	Approaches to studying benthic food supplies	44

Chapter 4	Behaviour, diet and energy intake	
	Abstract	47
	Introduction	48
	Methods	49
	Results	
	Roosting areas	52
	Feeding areas	52
	Distances travelled during feeding	54
	Site fidelity to roosting areas	54
	Activity	57
	Feeding method and prey type	61
	Pied Oystercatcher: diet and intake rates	63
	Red Knot: diet and intake rate	68
	Bar-tailed Godwit: diet and intake rates	76
	Discussion	
	Use of space	77
	Feeding time	79
	Pied Oystercatcher: energy requirements, diet and energy intake	81
	Red Knot: diet, energy requirements and energy intake	84
	Bar-tailed Godwit: diet, energy demand and energy intake	90
	Dietary overlap between species	92
	Seasonal energy demands and intake	92
Chapter 5	Seasonal mass changes in knots in New Zealand	
	Abstract	94
	Introduction	95
	Methods	95
	Results	96
	Discussion	99
	Is the juvenile mass increase adaptive?	101

Chapter 6	Body composition of Knots <i>Calidris canutus rogersi</i> from northern New Zealand in the premigratory period (Co-authored with Dr. Theunis Piersma, submitted to <i>Notornis</i>)	
	Abstract	105
	Introduction	106
	Methods	108
	Results	109
	Discussion	117
Chapter 7	The northward migration of arctic waders in New Zealand: departure behaviour, timing and possible migration routes of Red Knots and Bar-tailed Godwits from Farewell Spit, North-West Nelson (Submitted to <i>Emu</i>)	
	Abstract	121
	Introduction	122
	Methods	124
	Results	
	Premigratory and migratory behaviour	126
	Patterns of departures	128
	Direction of departures	130
	Influence of weather	132
	How far might New Zealand knots be able to fly?	135
	Discussion	
	Migratory behaviour	135
	Timing of departure	136
	Effect of weather	139
	Flight ranges	139
	Do knots fly to Australia direct or via Northland?	140

Chapter 8	Population processes and wader limitation: a theoretical excursion	143
	Carrying capacity: concepts and utility	144
	Mortality and selection	151
Chapter 9	General discussion	
	Food supply and usage	154
	Migratory preparation	155
	Numbers of birds on Farewell Spit: are they limited?	156
	Evidence for limitation?	161
	Interactions with Black Swans	163
	Energetic costs: implications for wader distributions around New Zealand	164
	Future research	167
	References	170

List of Figures and Tables

Figures

- Figure 2.1** Location of Farewell Spit
- Figure 2.2** Detail of study site at the tip of Farewell Spit
- Figure 2.3** Mean windspeed recorded at the Farewell Spit lighthouse.
- Figure 2.4** Mean temperature at the Farewell Spit lighthouse.
-
- Figure 3.1** How the harvestability of a prey item depends on its ingestibility, profitability and accessibility.
- Figure 3.2** Biomass of main taxonomic groups of organisms at Tip1 sampling site.
- Figure 3.3** Biomass of main taxonomic groups of organisms at Tip2 sampling site.
- Figure 3.4** Biomass of main taxonomic groups of organisms at Pan and Tip1-Tip2 sampling sites.
- Figure 3.5** Biomass of main taxonomic groups of organisms at Flat1 and Chan1 sampling sites.
- Figure 3.6** Pied Oystercatcher prey biomass levels at four main sampling sites.
- Figure 3.7** Knot prey biomass levels at three main sampling sites.
- Figure 3.8** Godwit prey biomass levels at four main sampling sites.
- Figure 3.9** Relationship between hinge height and shell length for cockles.
- Figure 3.10** Relationship between hinge+top height and shell length in cockles. *
- Figure 3.11** Ingestibility limitation of cockles for knots.
- Figure 3.12** Relationship between hinge+top length and shell length in the pipi.
- Figure 3.13** Ingestibility limit for knots feeding in pipi.
- Figure 3.14** Mean cockle shell length and mean deviation by month (1994).
- Figure 3.15** Size distribution of cockles at Tip3 in January, June and December 1994.
- Figure 3.16** Size distributions of mussels present near the tip of Farewell Spit in late summer (February) and early winter (June).
- Figure 3.17** Seasonal changes in flesh content of 15, 25 and 35 mm cockles.
- Figure 3.18** Seasonal changes in density of cockles at Tip3 in 1994.
- Figure 3.19** Seasonal changes in total biomass in cockles at Tip3 in 1994.
- Figure 3.20** Seasonal changes in flesh content in *Macomona* of 15, 25 and 35 mm length in 1994.
- Figure 3.21** Average depth of 15, 25 and 35 mm *Macomona*.
- Figure 3.22** Seasonal variation in harvestable biomass of *Macomona* to knots and oystercatchers.

- Figure 3.23** Flesh mass as a function of shell length in pipi over summer (November-April).
- Figure 3.24** Ratio of predicted summer mass to winter mass for four sizes of pipi.
- Figure 3.25** Flesh content as a function of shell length in mussels in late summer
- Figure 4.1.** Generalised movements of feeding waders at low tide.
- Figure 4.2** Counts of knots roosting near Bush end Point and the shellbanks from 14-1-1994 to 8-3-1994.
- Figure 4.3** Counts of godwits roosting near Bush end Point and the shellbanks from 14-1-1994 to 8-3-1994.
- Figure 4.4** Percentage of feeding knots over all tides in the basic period, relative to tide.
- Figure 4.5** Percentage of feeding godwits over all tides in the basic period, relative to tide.
- Figure 4.6** Percentage of feeding knots over all tides in the premigratory period, relative to tide.
- Figure 4.7** Percentage of feeding godwits over all tides in the premigratory period, relative to tide.
- Figure 4.8** Percentage of feeding oystercatchers over all tides in summer, relative to tide.
- Figure 4.9** Percentage of feeding oystercatchers over all tides in winter, relative to tide.
- Figure 4.10** Size of mussels eaten by Pied Oystercatchers at the hummocks in February and June 1994.
- Figure 4.11** Mussel selection by oystercatchers at the hummocks.
- Figure 4.12** Proportion of biomass per mm size class contributing to diet of oystercatchers feeding on mussels.
- Figure 4.13** Estimated daily energy intake of Pied Oystercatchers feeding on four prey types in summer and early winter.
- Figure 4.14** Localities used in grouping of knot faeces
- Figure 4.15** Size distribution of pipi recovered from knot faeces (all localities combined).
- Figure 4.16** Size distribution of pipi recovered from knot faeces and present in sediment for 1993/94 over all sites.
- Figure 4.17** Size distributions of pipi taken by knots, estimated from faeces and regurgitates.
- Figure 4.18** Size distribution of cockles taken by knots with all localities combined, estimated from faeces.
- Figure 4.19** Size distribution of cockles from knot faeces and regurgitates in 1993/94.
- Figure 4.20** Size distribution of cockles taken by knots (estimated from faeces) and present in sediment for 1993/94 over all sites.
- Figure 4.21** Intake rate of knots over all sites and times.
- Figure 4.22** Maintenance metabolism of Pied Oystercatchers at Farewell Spit.
- Figure 4.23** Maintenance metabolism of Red Knots on Farewell Spit.

- Figure 4.24** Maintenance metabolism of Bar-tailed Godwits on Farewell Spit.
- Figure 5.1** Localities of knot catching operations in the Auckland region.
- Figure 5.2** Mass of knots though the year, separated into 3rd, 2nd and 1st-year birds.
- Figure 6.1** Schematic representation of how a sample of birds on a single day can approximate the "fattening" trajectory of a hypothetical individual.
- Figure 6.2** Relationship between body mass and total fat mass and total fat-free dry mass for knots.
- Figure 6.3** Relationship between the four components of total fat mass, and total fat mass in knots.
- Figure 6.4** Total FFDM plotted against fat mass in knots.
- Figure 6.5** Organ FFDM plotted against total fat mass in knots.
- Figure 7.1** Map of localities in Australasia mentioned in the text.
- Figure 7.2** Daily pattern of observed departures of knots and godwits from Farewell Spit, March 1994.
- Figure 7.3** Pattern of departures relative to time of day (top), and observer time in field (lower).
- Figure 7.4** Pattern of observed departures relative to tidal stage.
- Figure 7.5** Directions of departures for knot and godwit.
- Figure 7.6** Degree of wind drift experienced by departing flocks.
- Figure 7.7** Wind vectors for departing flocks.
- Figure 7.8** Plot of departure occurrences relative to prevailing weather systems.
- Figure 7.9** Estimated flight range of knots.
- Figure 7.10** Tidal Influence Index, plotted against tidal height.
- Figure 8.1** Effects of habitat loss on bird populations.
- Figure 9.1** Count of knots at Farewell Spit from 1983 to 1995.
- Figure 9.2** Counts of godwits at Farewell Spit from 1983 to 1995.
- Figure 9.3** Percentage deviation from three-year running mean for counts of knots on Farewell Spit
- Figure 9.4** Percentage deviation from three-year running mean for counts of godwits on Farewell Spit.
- Figure 9.5** Count of oystercatchers at Farewell Spit from 1984 to 1995.

- Figure 9.6** Percentage deviation from three-year running mean for winter counts of oystercatchers on Farewell Spit
- Figure 9.7** Maintenance metabolism for knots at the Firth of Thames, Farewell Spit and Southland.
- Figure 9.8** Maintenance metabolism for knots at the Firth of Thames, Farewell Spit and Southland.

Tables

- Table 3.1** Sampling dates for invertebrates.
- Table 3.2** Intercept, exponent and explained variance for power functions relating shell length to biomass (mg AFDM) for cockles in different months.
- Table 3.3** Intercept, exponent and explained variance for power functions relating shell length to biomass (mg AFDM) for *Macomona* in different months.
- Table 4.1** Feeding method on main prey types for knots, godwits and oystercatchers.
- Table 4.2** Intake, pecking/probing and pacing rates of foraging oystercatchers.
- Table 4.3** Composition of knot diet assessed from faeces according to locality, for all years combined.
- Table 4.4** Mean size (mm) of pipi taken by knots, summarised by locality and summer.
- Table 4.5** Mean size (mm) of cockles taken by knots, summarised by locality and summer.
- Table 4.6** Energy intake of godwits feeding on worms (*Travisia* and 'slender worms').
- Table 6.1** Physical and plumage details of all analysed knots, listed in order of decreasing total body mass.
- Table 6.2** Correlations of body mass, fat mass, total FFDM, FFDM of all body parts and three structural measures, with fat mass and total FFDM for adult knots.
- Table 6.3** Comparison of body part masses for adult and juvenile knots.
- Table 7.1** Flock sizes, vocalisation and presence of visible sun at departures for knot and godwit, at Farewell Spit, March 1994.
- Table 7.2** Departing flock formation and changes in knots and godwits at Farewell Spit, March 1994.